

EIGENVALUES OF THE LAPLACIAN ON A COMPACT MANIFOLD WITH DENSITY

BRUNO COLBOIS, AHMAD EL SOUFI, AND ALESSANDRO SAVO

ABSTRACT. In this paper, we study the spectrum of the weighted Laplacian (also called Bakry-Emery or Witten Laplacian) L_σ on a compact, connected, smooth Riemannian manifold (M, g) endowed with a measure σdv_g . First, we obtain upper bounds for the k -th eigenvalue of L_σ which are consistent with the power of k in Weyl's formula. These bounds depend on integral norms of the density σ , and in the second part of the article, we give examples showing that this dependence is, in some sense, sharp. As a corollary, we get bounds for the eigenvalues of Laplace type operators, such as the Schrödinger operator or the Hodge Laplacian on p -forms. In the special case of the weighted Laplacian on the sphere, we get a sharp inequality for the first nonzero eigenvalue which extends Hersch's inequality.

1. INTRODUCTION

In this article, our main aim is to study the spectrum of the weighted Laplacian (also called Bakry-Emery Laplacian) L_σ on a compact, connected, smooth Riemannian manifold (M, g) endowed with a measure σdv_g , where $\sigma = e^{-f} \in C^2(M)$ is a positive density and dv_g is the Riemannian measure induced by the metric g . Such a triple (M, g, σ) is known in literature as a weighted Riemannian manifold, a manifold with density, a smooth metric measure space or a Bakry-Emery manifold. Denoting by ∇^g and Δ_g the gradient and the Laplacian with respect to the metric g , the operator L_σ is defined by

$$L_\sigma = \Delta_g - \frac{1}{\sigma} \nabla^g \sigma \cdot \nabla^g = \Delta_g + \nabla^g f \cdot \nabla^g$$

so that, for any function $u \in C^2(M)$, satisfying Neumann boundary conditions if $\partial M \neq \emptyset$,

$$\int_M |\nabla^g u|^2 \sigma dv_g = \int_M u L_\sigma u \sigma dv_g.$$

Note that L_σ is self-adjoint as an operator on $L^2(\sigma dv_g)$ and is unitarily equivalent (through the transform $\sqrt{\sigma} : L^2(\sigma dv_g) \rightarrow L^2(dv_g)$) to the Schrödinger operator $H_\sigma = \Delta_g + \frac{1}{4} |\nabla^g f|^2 + \frac{1}{2} \Delta_g f$, which is nothing but the restriction to functions of the Witten Laplacian associated to f . That is why L_σ itself is sometimes called *Witten Laplacian*.

Weighted manifolds arise naturally in several situations in the context of geometric analysis and their study has been very active in recent years. Their

2010 *Mathematics Subject Classification.* 58J50, 35P15, 47A75.

Key words and phrases. Manifold with density, Weighted Laplacian, Witten Laplacian, Eigenvalue, Upper bound.

Bakry-Emery curvature $\text{Ric}_\sigma = \text{Ric}_g + \text{Hess}f$ plays a role which is similar in many respects to that played by the Ricci curvature for Riemannian manifolds, and appears as a centerpiece in the analysis of singularities of the Ricci flow in Perelman's work (see [35, 36]). The weighted Laplacian L_σ appears naturally in the study of diffusion processes (see e.g., the pioneering work of Bakry and Emery [2]). Eigenvalues of L_σ are strongly related to asymptotic properties of mm-spaces, such as the study of Levy families (see [11, 19, 34]). Without being exhaustive, we refer to the following articles and the references therein: [29, 30, 31, 37, 38, 39, 43] and, closely related to our topic, [1, 12, 13, 21, 32, 33, 41, 44, 45]

The spectrum of L_σ , with Neumann boundary conditions if $\partial M \neq \emptyset$, consists of an unbounded sequence of eigenvalues

$$\text{Spec}(L_\sigma) = \{0 = \lambda_1(L_\sigma) < \lambda_2(L_\sigma) \leq \lambda_3(L_\sigma) \leq \dots \leq \lambda_k(L_\sigma) \leq \dots\}$$

which satisfies the Weyl's asymptotic formula

$$\lambda_k(L_\sigma) \sim 4\pi^2 \omega_n^{-\frac{2}{n}} \left(\frac{k}{V_g(M)} \right)^{\frac{2}{n}}, \quad \text{as } k \rightarrow \infty$$

where $V_g(M)$ is the Riemannian volume of (M, g) and ω_n is the volume of the unit ball in \mathbb{R}^n . The first aim of this paper is to obtain bounds for $\lambda_k(L_\sigma)$ which are consistent with the power of k in Weyl's formula.

Before stating our results, let us recall some known facts about the eigenvalues $(\lambda_k(g))_{k \geq 1}$ of the usual Laplacian Δ_g (case $\sigma = 1$). Firstly, the well-known Hersch's isoperimetric inequality (see [24]) asserts that on the 2-dimensional sphere \mathbb{S}^2 , the first positive normalized eigenvalue $\lambda_2(g)V_g(M)$ is maximal when g is a "round" metric (see [9, 10, 25, 27, 40] for similar results on other surfaces). Korevaar [26] proved that on any compact manifold of dimension $n = 2$, $\lambda_k(g)V_g(M)$ is bounded above independently of g . More precisely, if M is a compact orientable surface of genus γ , then

$$\lambda_k(g)V_g(M) \leq C(\gamma + 1)k \tag{1}$$

where C is an absolute constant (see [20] for an improved version of this inequality). On the other hand, on any compact manifold M of dimension $n \geq 3$, the normalized first positive eigenvalue $\lambda_2(g)V_g(M)^{2/n}$ can be made arbitrarily large when g runs over the set of all Riemannian metrics on M (see [4, 28]). However, the situation changes as soon as we restrict ourselves to a fixed conformal class of metrics. Indeed, on the sphere \mathbb{S}^n , round metrics maximize $\lambda_2(g)V_g(\mathbb{S}^n)^{2/n}$ among all metrics g which are conformally equivalent to the standard one (see [8, Proposition 3.1]). Furthermore, Korevaar [26] proved that for any compact Riemannian manifold (M, g) one has

$$\lambda_k(g)V_g(M)^{2/n} \leq C([g])k^{2/n} \tag{2}$$

where $C([g])$ is a constant depending only on the conformal class $[g]$ of the metric g . Korevaar's approach has been revisited and placed in the context of metric measure spaces by Grigor'yan and Yau [16] and, then, by Grigor'yan, Netrusov and Yau [18].

The first observation we can make about possible extensions of these results to weighted Laplacians is that, given any compact Riemannian manifold (M, g) , the eigenvalues $\lambda_k(L_\sigma)$ cannot be bounded above independently

of σ . Indeed, from the semi-classical analysis of the Witten Laplacian (see [22]), we can easily deduce that (Proposition 2.1) if f is any smooth Morse function on M with m_0 stable critical points, then the family of densities $\sigma_\varepsilon = e^{-f/\varepsilon}$ satisfies for $k > m_0$,

$$\lambda_k(L_{\sigma_\varepsilon}) \xrightarrow{\varepsilon \rightarrow 0} +\infty.$$

Therefore, any extension of the inequalities (1) and (2) to L_σ must necessarily have a density dependence in the right-hand side. The following theorem gives such an extension in which the upper bound depends on the ratio between the $L^{\frac{n}{n-2}}$ -norm and the L^1 -norm of σ . In all the sequel, the L^p norm of σ with respect to dv_g will be denoted by $\|\sigma\|_p^g$.

Theorem 1.1. *Let (M, g, σ) be a compact weighted Riemannian manifold. The eigenvalues of the operator L_σ , with Neumann boundary conditions if $\partial M \neq \emptyset$, satisfy :*

(I) *If $n \geq 3$, then, $\forall k \geq 1$,*

$$\lambda_k(L_\sigma) \leq C([g]) \frac{\|\sigma\|_{\frac{n}{n-2}}^g}{\|\sigma\|_1^g} k^{2/n}$$

where $C([g])$ is a constant depending only on the conformal class of g .

(II) *if $n = 2$ and M is orientable of genus γ , then, $\forall k \geq 1$,*

$$\lambda_k(L_\sigma) \leq C \frac{\|\sigma\|_\infty}{\|\sigma\|_1^g} (\gamma + 1) k$$

where C is an absolute constant.

It is clear that taking $\sigma = 1$ in Theorem 1.1, we recover the inequalities (1) and (2). Moreover, as we will see in the next section, if M is boundaryless and the conformal class $[g]$ contains a metric g_0 with nonnegative Ricci curvature, then $C([g]) \leq C(n)$, where $C(n)$ is a constant which depends only on the dimension.

Notice that there already exist upper bounds for the eigenvalues of L_σ in the literature, but they usually depend on derivatives of σ , either directly or indirectly, through the Bakry-Emery curvature. The main feature of our result is that the upper bounds we obtain depend only on L^p -norms of the density. The proof of Theorem 1.1 relies in an essential way on the technique developed by Grigor'yan, Netrusov and Yau [18].

Regarding Hersch's isoperimetric type inequalities, they extend to our context as follows (see Corollary 3.2): Given any metric g on \mathbb{S}^n which is conformally equivalent to the standard metric g_0 , and any positive density $\sigma \in C^2(\mathbb{S}^n)$, one has

$$\lambda_2(L_\sigma) \leq n |\mathbb{S}^n|^{\frac{n}{2}} \frac{\|\sigma\|_{\frac{n}{n-2}}^g}{\|\sigma\|_1^g}$$

where $|\mathbb{S}^n|$ is the volume of the standard n -sphere and with the convention that $\|\sigma\|_{\frac{n}{n-2}}^g = \|\sigma\|_\infty$ when $n = 2$. Moreover, the equality holds in the inequality if and only if σ is constant and g is a round metric.

Next, let us consider a Riemannian vector bundle E over a Riemannian manifold (M, g) and a Laplace type operator

$$H = D^*D + T$$

acting on smooth sections of the bundle. Here D is a connection on E which is compatible with the Riemannian metric and T is a symmetric bundle endomorphism (see e.g., [3, Section E]). The operator H is self-adjoint and elliptic and we will list its eigenvalues as: $\lambda_1(H) \leq \lambda_2(H) \leq \dots \leq \lambda_k(H) \leq \dots$. Important examples of such operators are given by Schrödinger operators acting on functions (here T is just the potential), the Hodge Laplacian acting on differential forms (in which case T is the curvature term in Bochner's formula), and the square of the Dirac operator (T being in this case a multiple of the scalar curvature). Another important example is the Witten Laplacian acting on differential forms, whose restriction to functions is precisely given by a weighted Laplacian, the main object of study of this paper.

In Section 4 we will prove (Theorem 4.1) an upper bound for the gap between the k -th eigenvalue and the first eigenvalue of H involving integral norms of a first eigensection ψ . For example, if $n \geq 3$, then

$$\lambda_k(H) - \lambda_1(H) \leq C([g]) \left(\frac{\|\psi\|_{\frac{2n}{n-2}}^g}{\|\psi\|_2^g} \right)^2 k^{2/n}, \quad (3)$$

where $C[g]$ is a constant depending only on the conformal class of g . The reason why we bound the gap instead of $\lambda_k(H)$ itself is due to the fact that, even when $\sigma = 1$ and H is the standard Hodge Laplacian acting on p -forms, the first positive eigenvalue is not bounded on any conformal class of metrics (see [5]). For estimates on the gap when a finite group of isometries is acting, we refer to [7].

Inequality (3) should be regarded as an extension of Theorem 1.1. Indeed, if $H_\sigma = \sqrt{\sigma} L_\sigma \frac{1}{\sqrt{\sigma}}$ is the Schrödinger operator which is unitarily equivalent to the operator L_σ , then $\lambda_1(H_\sigma) = \lambda_1(L_\sigma) = 0$ and any first eigenfunction of H_σ is a scalar multiple of $\sqrt{\sigma}$. Thus, taking $\psi = \sqrt{\sigma}$ in (3) we recover the first estimate in Theorem 1.1.

Our main aim in section 5 is to discuss the accuracy of the upper bounds given in Theorem 1.1 regarding the way they depend on the density σ . That is why we exhibit an explicit family of compact manifolds (M, g) , each endowed with a sequence of densities $\{\sigma_j\}$, and give a sharp lower estimate of the first positive eigenvalue $\lambda_2(L_{\sigma_j})$ in terms of j . This enables us to see that both $\lambda_2(L_{\sigma_j})$ and the ratio $\|\sigma_j\|_{\frac{n}{n-2}}^g / \|\sigma_j\|_1^g$ tend to infinity linearly with respect to j . Thus, we have

$$A \frac{\|\sigma_j\|_{\frac{n}{n-2}}^g}{\|\sigma_j\|_1^g} \leq \lambda_2(L_{\sigma_j}) \leq B \frac{\|\sigma_j\|_{\frac{n}{n-2}}^g}{\|\sigma_j\|_1^g}$$

with $\lambda_2(L_{\sigma_j}) \rightarrow +\infty$ as $j \rightarrow +\infty$, and A and B are two positive constants which do not depend on j .

The examples of densities we give are modeled on Gaussian densities (i.e. $\sigma_j(x) = e^{-j|x|^2}$) on \mathbb{R}^n . For example, if Ω is a bounded convex domain in

\mathbb{R}^n , we observe that $\lambda_2(L_{\sigma_j}) \geq 2j$ for all j . We then extend the lower bound to manifolds of revolution (at least asymptotically as $j \rightarrow \infty$). However, in the case of a closed manifold of revolution, there is an additional difficulty coming from the fact that we need to extend smoothly this kind of density to the whole manifold in such a way as to preserve the estimates on both the eigenvalues and the L^p -norms.

2. UPPER BOUNDS FOR WEIGHTED EIGENVALUES IN A SMOOTH METRIC MEASURE SPACE AND PROOF OF THEOREM 1.1

Let (M, g) be a compact connected Riemannian manifold, possibly with a non-empty boundary. Let $\sigma \in L^\infty(M)$ be a bounded nonnegative function on M and let ν be a non-atomic Radon measure on M with $0 < \nu(M) < \infty$. To such a pair (σ, ν) , we associate the sequence of non-negative numbers $\{\mu_k(\sigma, \nu)\}_{k \in \mathbb{N}}$ given by

$$\mu_k(\sigma, \nu) = \inf_{E \in S_k} \sup_{u \in E \setminus \{0\}} R_{\sigma, \nu}(u)$$

where S_k is the set of all k -dimensional vector subspaces of $H^1(M)$ and

$$R_{\sigma, \nu}(u) = \frac{\int_M |\nabla^g u|_g^2 \sigma dv_g}{\int_M u^2 d\nu}.$$

In the case where σ is of class C^2 and $\nu = \sigma dv_g$, the variational characterization of eigenvalues of the weighted Laplacian $L_\sigma = \Delta_g - \frac{1}{\sigma} \nabla^g \sigma \cdot \nabla^g$ gives (see e.g. [17])

$$\lambda_k(L_\sigma) = \mu_k(\sigma, \sigma dv_g). \quad (4)$$

Theorem 1.1 is a direct consequence of the following

Theorem 2.1. *Let (M, g) be a compact Riemannian manifold possibly with nonempty boundary. Let $\sigma \in L^\infty(M)$ be a nonnegative function and let ν be a non-atomic Radon measure with $0 < \nu(M) < \infty$.*

(I) *If $n \geq 3$, then for every $k \geq 1$, we have*

$$\mu_k(\sigma, \nu) \leq C([g]) \frac{\|\sigma\|_{\frac{n}{n-2}}^g}{\nu(M)} k^{2/n},$$

where $C([g])$ is a constant depending only on the conformal class of g . Moreover, if M is closed and $[g]$ contains a metric with nonpositive Ricci curvature, then $C([g]) \leq C(n)$ where $C(n)$ is a constant depending only on n .

(II) *If M is a compact orientable surface of genus γ , then for every $k \geq 1$, we have*

$$\mu_k(\sigma, \nu) \leq C \frac{\|\sigma\|_\infty}{\nu(M)} (\gamma + 1) k.$$

where C is an absolute constant.

The proof of this theorem is based on the method described by Grigor'yan, Netrusov and Yau in [18] and follows the same lines as the proof they have given in the case $\sigma = 1$. The main step consists in the construction of a family of disjointly supported functions with controlled Rayleigh quotient.

Let us fix a reference metric $g_0 \in [g]$ and denote by d_0 the distance associated to g_0 . An *annulus* $A \subset M$ is a subset of M of the form $\{x \in M : r < d_0(x, a) < R\}$ where $a \in M$ and $0 \leq r < R$ (if necessary, we will denote

it $A(a, r, R)$). The annulus $2A$ is by definition the annulus $\{x \in M : r/2 < d_0(x, a) < 2R\}$.

To such an annulus we associate the function u_A supported in $2A$ and such that

$$u_A(x) = \begin{cases} 1 - \frac{2}{r}d_0(x, A) & \text{if } \frac{r}{2} \leq d_0(x, a) \leq r \\ 1 & \text{if } x \in A \\ 1 - \frac{1}{R}d_0(x, A) & \text{if } R \leq d_0(x, a) \leq 2R \end{cases}$$

We introduce the following constant:

$$\Gamma(g_0) = \sup_{x \in M, r > 0} \frac{V_{g_0}(B(x, r))}{r^n}$$

where $B(x, r)$ stands for the ball of radius r centered at x in (M, d_0) . Notice that since M is compact, the constant $\Gamma(g_0)$ is finite and depends only on g_0 . This constant can be bounded from above in terms of a lower bound of the Ricci curvature Ric_{g_0} and an upper bound of the diameter $diam(M, g_0)$ (Bishop-Gromov inequality). In particular, if the Ricci curvature of g_0 is nonnegative, then $\Gamma(g_0)$ is bounded above by a constant depending only on the dimension n .

Lemma 2.1. *For every annulus $A \subset (M, d_0)$ one has*

$$\int_M |\nabla^g u_A|^2 \sigma dv_g \leq 8 \Gamma(g_0)^{\frac{2}{n}} \left(\int_{2A} \sigma^{\frac{n}{n-2}} dv_g \right)^{1-\frac{2}{n}}.$$

Proof. Let $A = A(a, r, R)$ be an annulus of (M, d_0) . Since u_A is supported in $2A$ we get, using Hölder inequality,

$$\int_M |\nabla^g u_A|^2 \sigma dv_g = \int_{2A} |\nabla^g u_A|^2 \sigma dv_g \leq \left(\int_{2A} |\nabla^g u_A|^n dv_g \right)^{\frac{2}{n}} \left(\int_{2A} \sigma^{\frac{n}{n-2}} dv_g \right)^{1-\frac{2}{n}}.$$

From the conformal invariance of $\int_{2A} |\nabla^g u_A|^n dv_g$ we have

$$\int_{2A} |\nabla^g u_A|^n dv_g = \int_{2A} |\nabla^{g_0} u_A|^n dv_{g_0}$$

with

$$|\nabla^{g_0} u_A| \stackrel{a.e.}{=} \begin{cases} \frac{2}{r} & \text{if } \frac{r}{2} \leq d_0(x, a) \leq r \\ 0 & \text{if } r \leq d_0(x, a) \leq R \\ \frac{1}{R} & \text{if } R \leq d_0(x, a) \leq 2R. \end{cases}$$

Hence,

$$\int_{2A} |\nabla^{g_0} u_A|^n dv_{g_0} \leq \left(\frac{2}{r} \right)^n V_{g_0}(B(a, r)) + \left(\frac{1}{R} \right)^n V_{g_0}(B(a, 2R)) \leq 2^{n+1} \Gamma(g_0)$$

where the last inequality follows from the definition of $\Gamma(g_0)$. Putting together all the previous inequalities, we obtain the result of the Lemma. \square

Proof of part (I) of Theorem 2.1: Let us introduce the constant $N(M, d_0)$, that we call the *covering constant*, defined to be the infimum of the set of all integers N such that, for all $r > 0$, any ball of radius $2r$ in (M, d_0) can be covered by N balls of radius r . Again, the compactness of M ensures that $N(M, d_0)$ is finite, and Bishop-Gromov inequality allows us to bound it from above in terms of the dimension when the Ricci curvature of (M, g_0) is nonnegative.

Since the metric measure space (M, d_0, ν) has a finite covering constant and a non atomic measure, one can apply Theorem 1.1 of [18] and conclude that there exists a constant $c(N)$ depending only on $N(M, d_0)$ such that for each positive integer k , there exists a family of $2k$ annuli A_1, \dots, A_{2k} on M such that the annuli $2A_1, \dots, 2A_{2k}$ are mutually disjoint and, $\forall i \leq 2k$,

$$\nu(A_i) \geq c(N) \frac{\nu(M)}{k}. \quad (5)$$

Since the annuli $2A_i$ are mutually disjoint, one has

$$\sum_{i \leq 2k} \int_{2A_i} \sigma^{\frac{n}{n-2}} dv_g \leq \int_M \sigma^{\frac{n}{n-2}} dv_g.$$

Thus, at most k annuli among $2A_1, \dots, 2A_{2k}$ satisfy $\int_{2A_i} \sigma^{\frac{n}{n-2}} dv_g > \frac{1}{k} \int_M \sigma^{\frac{n}{n-2}} dv_g$. Therefore, we can assume without loss of generality that the k annuli A_1, \dots, A_k satisfy

$$\left(\int_{2A_i} \sigma^{\frac{n}{n-2}} dv_g \right)^{1-\frac{2}{n}} \leq \frac{1}{k^{1-\frac{2}{n}}} \|\sigma\|_{\frac{n}{n-2}}^g.$$

The corresponding functions u_{A_1}, \dots, u_{A_k} are such that (Lemma 2.1)

$$\begin{aligned} R_{\sigma, \nu}(u_{A_i}) &= \frac{\int_M |\nabla^g u_{A_i}|_g^2 \sigma dv_g}{\int_M u_{A_i}^2 d\nu} \leq \frac{1}{\nu(A_i)} 8\Gamma(g_0)^{\frac{2}{n}} \left(\int_{2A_i} \sigma^{\frac{n}{n-2}} dv_g \right)^{1-\frac{2}{n}} \\ &\leq \frac{1}{\nu(A_i)} 8\Gamma(g_0)^{\frac{2}{n}} \frac{\|\sigma\|_{\frac{n}{n-2}}^g}{k^{1-\frac{2}{n}}}. \end{aligned}$$

Using inequality (5), we get

$$R_{\sigma, \nu}(u_{A_i}) \leq 8 \frac{\Gamma(g_0)^{\frac{2}{n}}}{c(N(M, d_0))} \frac{\|\sigma\|_{\frac{n}{n-2}}^g}{\nu(M)} k^{\frac{2}{n}}.$$

Since the functions u_{A_1}, \dots, u_{A_k} are disjointly supported, they form a k -dimensional subspace on which the Rayleigh quotient is bounded above by the right hand side of the last inequality. We set $C([g]) = 8 \frac{\Gamma(g_0)^{\frac{2}{n}}}{c(N(M, d_0))}$ and conclude using the min-max formula.

As we mentioned above, if M is closed and the Ricci curvature of g_0 is non negative, then the constants $\Gamma(g_0)$, $N(M, d_0)$ and, hence, $C([g])$ are bounded in terms of the dimension n .

Proof of part (II) of Theorem 2.1. Assume now that (M, g) is a compact orientable surface of genus γ , possibly with boundary, and let $\rho \in C^\infty(M)$ be a positive function on M . If M has nonempty boundary, then we glue a disk on each boundary component of M and extend ν by 0. This closed surface admits a conformal branched cover ψ over \mathbb{S}^2 with degree $\deg(\psi) \leq \lfloor \frac{\gamma+3}{2} \rfloor$. We endow \mathbb{S}^2 with the usual spherical distance d_0 and the pushforward measure $\mu = \psi_*(\nu)$. We apply Theorem 1.1 of [18] to the metric measure space (\mathbb{S}^2, d_0, μ) and deduce that there exist an absolute constant $c = c(N(\mathbb{S}^2, d_0))$ and k annuli $A_1, \dots, A_k \subset \mathbb{S}^2$ such that the annuli $2A_1, \dots, 2A_k$ are mutually disjoint and, $\forall i \leq k$,

$$\mu(A_i) \geq c \frac{\mu(\mathbb{S}^2)}{k}. \quad (6)$$

We set for each $i \leq k$, $v_i = u_{A_i} \circ \psi$. From the conformal invariance of the energy and Lemma 2.1, one deduces that, for every $i \leq k$,

$$\int_M |\nabla_g v_i|^2 dv_g = \deg(\psi) \int_{\mathbb{S}^2} |\nabla_{g_0} u_{A_i}|^2 dv_{g_0} \leq 8\Gamma(\mathbb{S}^2, g_0) \lfloor \frac{\gamma+3}{2} \rfloor,$$

while, since u_{A_i} is equal to 1 on A_i ,

$$\int_M v_i^2 d\nu \geq \nu(\psi^{-1}(A_i)) = \mu(A_i) \geq c \frac{\mu(\mathbb{S}^2)}{k} = c \frac{\nu(M)}{k}.$$

Therefore,

$$R_{\sigma, \nu}(v_i) \leq \frac{8\Gamma(\mathbb{S}^2, g_0)}{c\nu(M)} \lfloor \frac{\gamma+3}{2} \rfloor k \leq \frac{C(\gamma+1)}{\nu(M)} k$$

where C is an absolute constant. Noting that the k functions v_1, \dots, v_k are disjointly supported in M , we deduce the desired inequality for $\mu_k(\sigma, \nu)$.

We end this section with the following observation showing that the presence of the density in the RHS of the inequalities of Theorem 1.1 is essential. This question will also be discussed in Section 5.

Proposition 2.1. *Let (M, g) be a compact Riemannian manifold and let f be a smooth Morse function on M . For every $\varepsilon > 0$ we set $\sigma_\varepsilon = e^{-f/\varepsilon}$. If m_0 denotes the number of stable critical points of f , then there exists $\varepsilon_0 > 0$ such that, $\forall \varepsilon \in (0, \varepsilon_0)$,*

$$\lambda_{m_0+1}(L_{\sigma_\varepsilon}) \geq \frac{1}{\sqrt{\varepsilon}}.$$

Proof. First observe that for any density $\sigma = e^{-f}$, the operator L_σ is unitarily equivalent to the Schrödinger operator

$$H_\sigma = \Delta_g - \frac{\Delta_g \sqrt{\sigma}}{\sqrt{\sigma}} = \Delta_g + \frac{1}{4} |\nabla^g f|^2 + \frac{1}{2} \Delta_g f$$

acting on $L^2(dv_g)$ (indeed, $L_\sigma = \frac{1}{\sqrt{\sigma}} H_\sigma \sqrt{\sigma}$, where multiplication by $\sqrt{\sigma}$ is a unitary transform from $L^2(\sigma dv_g)$ to $L^2(dv_g)$). Consequently, denoting by $(\lambda_k(\varepsilon))_{k \geq 1}$ the eigenvalues of the semiclassical Schrödinger operator $\varepsilon^2 \Delta_g + \frac{1}{4} |\nabla^g f|^2 + \frac{\varepsilon}{2} \Delta_g f$, we get

$$\lambda_k(L_{\sigma_\varepsilon}) = \frac{1}{\varepsilon^2} \lambda_k(\varepsilon).$$

According to [22] (see also [23, proposition 2.2]), there exists $\varepsilon_0 > 0$ such that, $\forall \varepsilon < \varepsilon_0$, the number of eigenvalues $\lambda_k(\varepsilon)$ contained in the interval $[0, \varepsilon^{3/2})$ is exactly m_0 , that is, $\lambda_{m_0}(\varepsilon) < \varepsilon^{3/2}$ and $\lambda_{m_0+1}(\varepsilon) \geq \varepsilon^{3/2}$. Therefore,

$$\lambda_{m_0+1}(L_{\sigma_\varepsilon}) = \frac{1}{\varepsilon^2} \lambda_{m_0+1}(\varepsilon) \geq \frac{1}{\sqrt{\varepsilon}}.$$

□

3. SHARP ESTIMATES FOR THE FIRST POSITIVE EIGENVALUE

Let (M, g) be a compact connected Riemannian n -dimensional manifold, possibly with a non-empty boundary. Li and Yau introduced in [27] the notion of conformal volume as follows : Given any immersion ϕ from M to the standard sphere $(\mathbb{S}^p, g_{\mathbb{S}^p})$ of dimension p , we denote by $V(\phi)$ the volume of M with respect to the metric $\phi^*g_{\mathbb{S}^p}$, and by $V_c(\phi)$ the supremum of $V(\gamma \circ \phi)$ as γ runs over the group of conformal diffeomorphisms of $(\mathbb{S}^p, g_{\mathbb{S}^p})$. The conformal volume of $(M, [g])$ is

$$V_c(M, [g]) = \inf_{p \geq n} \inf \{V_c(\phi) : \phi \in \text{conf}((M, g), \mathbb{S}^p)\}$$

where $\text{conf}((M, g), \mathbb{S}^p)$ is the set of all conformal immersions from (M, g) to \mathbb{S}^p .

With the same notations as in the previous section, we have the following

Theorem 3.1. *Let (M, g) be a compact Riemannian manifold possibly with nonempty boundary. Let $\sigma \in L^\infty(M)$ be a nonnegative function and let ν be a non-atomic Radon measure on M with $0 < \nu(M) < \infty$. One has*

$$\mu_2(\sigma, \nu) \leq nV_c(M, [g])^{2/n} \frac{\|\sigma\|_{\frac{n}{n-2}}^g}{\nu(M)}, \quad (7)$$

with the convention that $\|\sigma\|_{\frac{n}{n-2}} = \|\sigma\|_\infty$ if $n = 2$.

Proof. From the definition of $\mu_2(\sigma, \nu)$, it is clear that if $u \in C^1(M)$ is any nonzero function such that $\int_M u d\nu = 0$, then, taking for E the 2-dimensional vector space generated by constant functions and u , we have

$$\mu_2(\sigma, \nu) \leq \sup_{w \in E} R_{\sigma, \nu}(w) \leq R_{\sigma, \nu}(u).$$

Let $\phi \in \text{conf}((M, g), \mathbb{S}^p)$. Using standard center of mass lemma (see e.g., [15, Proposition 4.1.5]), there exists a conformal diffeomorphism γ of \mathbb{S}^p so that the Euclidean components of the map $\bar{\phi} = \gamma \circ \phi$ satisfy

$$\int_M \bar{\phi}_j d\nu = 0, \quad j \leq p+1.$$

Thus, for every $j \leq p+1$,

$$\mu_2(\sigma, \nu) \int_M \bar{\phi}_j^2 d\nu \leq \int_M |\nabla^g \bar{\phi}_j|^2 \sigma dv_g. \quad (8)$$

From the fact that ϕ is conformal one has $\bar{\phi}^*g_{\mathbb{S}^p} = \left(\frac{1}{n} \sum_{j \leq p+1} |\nabla^g \bar{\phi}_j|^2\right) g$ and

$$V(\bar{\phi}) = \int_M \left(\frac{1}{n} \sum_{j \leq p+1} |\nabla^g \bar{\phi}_j|^2\right)^{n/2} dv_g.$$

We sum up in (8) and use Hölder's inequality to get

$$\begin{aligned} \mu_2(\sigma, \nu) \nu(M) &\leq \int_M \sum_{j \leq p+1} |\nabla^g \bar{\phi}_j|^2 \sigma dv_g \\ &\leq nV(\bar{\phi})^{2/n} \|\sigma\|_{\frac{n}{n-2}}^g \leq nV_c(\phi)^{2/n} \|\sigma\|_{\frac{n}{n-2}}^g. \end{aligned}$$

The proof of the theorem follows immediately. \square

An immediate consequence of Theorem 3.1 is the following

Corollary 3.1. *Let (M, g) be a compact Riemannian manifold possibly with nonempty boundary and let $\sigma \in C^2(M)$ be a positive function. One has*

$$\lambda_2(L_\sigma) \leq nV_c(M, [g])^{2/n} \frac{\|\sigma\|_{\frac{n}{n-2}}^g}{\|\sigma\|_1^g}. \quad (9)$$

In [8], Ilias and the second author proved that if the Riemannian manifold (M, g) admits an isometric immersion into a Euclidean space \mathbb{R}^p whose Euclidean components are first non-constant eigenfunctions of the Laplacian Δ_g , then the following equality holds

$$\lambda_2(\Delta_g)V_g(M)^{2/n} = nV_c(M, [g])^{2/n}. \quad (10)$$

In this case, the inequality (9) reads

$$\lambda_2(L_\sigma) \leq \lambda_2(\Delta_g)V_g(M)^{2/n} \frac{\|\sigma\|_{\frac{n}{n-2}}^g}{\|\sigma\|_1^g} \quad (11)$$

where the equality holds whenever σ is a constant function. This proves the sharpness of the inequality of Corollary 3.1.

Notice that all compact rank one symmetric spaces satisfy (10) and hence (11). In particular, we have the following result that extends Hersch's isoperimetric inequality [24] and its generalization to higher dimensions [8].

Corollary 3.2. *Let g be a Riemannian metric on \mathbb{S}^n which is conformal to the standard metric $g_{\mathbb{S}^n}$, and let $\sigma \in C^2(\mathbb{S}^n)$ be a positive function. One has*

$$\lambda_2(L_\sigma) \leq n|\mathbb{S}^n|^{\frac{2}{n}} \frac{\|\sigma\|_{\frac{n}{n-2}}^g}{\|\sigma\|_1^g} \quad (12)$$

where $|\mathbb{S}^n|$ is the volume of the Euclidean unit sphere. Moreover, the equality holds in (12) if and only if σ is constant and g is homothetically equivalent to $g_{\mathbb{S}^n}$.

Proof. Let $\gamma = (\gamma_1, \dots, \gamma_{n+1}) : \mathbb{S}^n \rightarrow \mathbb{S}^n$ be conformal transformation of \mathbb{S}^n such that, for every $i \leq n+1$, we have

$$\int_{\mathbb{S}^n} \gamma_i \sigma \, dv_g = 0.$$

Using the same arguments as in the proof of Theorem 3.1, we get

$$\begin{aligned} \lambda_2(L_\sigma) \int_{\mathbb{S}^n} \gamma_i^2 \sigma \, dv_g &\leq \int_{\mathbb{S}^n} |\nabla^g \gamma_i|^2 \sigma \, dv_g, \quad i \leq n+1, \\ \lambda_2(L_\sigma) \|\sigma\|_1^g &\leq \int_{\mathbb{S}^n} \sum_{i \leq n+1} |\nabla^g \gamma_i|^2 \sigma \, dv_g \\ &\leq n \left(\int_{\mathbb{S}^n} \left(\frac{1}{n} \sum_{i \leq n+1} |\nabla^g \gamma_i|^2 \right)^{n/2} dv_g \right)^{2/n} \|\sigma\|_{\frac{n}{n-2}}^g \end{aligned}$$

with (since γ is conformal from (\mathbb{S}^n, g) to $(\mathbb{S}^n, g_{\mathbb{S}^n})$)

$$\int_{\mathbb{S}^n} \left(\frac{1}{n} \sum_{i \leq n+1} |\nabla^g \gamma_i|^2 \right)^{n/2} dv_g = V_{\gamma^* g_{\mathbb{S}^n}}(\mathbb{S}^n) = |\mathbb{S}^n|.$$

The inequality (12) follows immediately.

Assume that the equality holds in (12). This implies that

- $L_\sigma \gamma_i = \lambda_2(L_\sigma) \gamma_i$, $i \leq n+1$, and
- the function σ is constant if $n = 2$ and, if $n \geq 3$, the functions $\left(\sum_{i=1}^{n+1} |\nabla^g \gamma_i|^2 \right)^{n/2}$ and $\sigma^{\frac{n}{n-2}}$ are proportional.

An elementary computation gives

$$0 = \sum_{i \leq n+1} L_\sigma \gamma_i^2 = 2 \sum_{i \leq n+1} (\gamma_i L_\sigma \gamma_i - |\nabla^g \gamma_i|^2) = 2 \left(\lambda_2(L_\sigma) - \sum_{i \leq n+1} |\nabla^g \gamma_i|^2 \right).$$

From the previous facts we see that σ is constant in all cases and, since $\gamma^* g_{\mathbb{S}^n} = \left(\frac{1}{n} \sum_{j \leq n+1} |\nabla^g \gamma_j|^2 \right) g = \frac{\lambda_2(L_\sigma)}{n} g$, the metric g is homothetically equivalent to $g_{\mathbb{S}^n}$. \square

4. EIGENVALUES OF LAPLACE TYPE OPERATORS

In this section we show that Theorem 1.1 extends to a much more general framework to give upper bounds of the eigenvalues of certain operators acting on sections of vector bundles, precisely the Laplace-type operators defined in the introduction. Throughout this section, (M, g) denotes a compact Riemannian manifold without boundary. We will use the notations introduced in Section 2 and refer to [3, 14] for details on Laplace type operators.

Theorem 4.1. *Let $H = D^*D + T$ be an operator of Laplace type acting on sections of a Riemannian vector bundle E over (M, g) , and let ψ be an eigensection associated to $\lambda_1(H)$. One has for all $k \geq 1$*

$$\lambda_k(H) - \lambda_1(H) \leq \mu_k(\sigma, \nu)$$

with $\sigma = |\psi|^2$ and $\nu = |\psi|^2 dv_g$. Thus,

a) If $n \geq 3$ then

$$\lambda_k(H) - \lambda_1(H) \leq C([g]) \left(\frac{\|\psi\|_{\frac{2n}{n-2}}^g}{\|\psi\|_2^g} \right)^2 k^{2/n}.$$

where $C([g])$ is a constant depending only on the conformal class of g .

b) If M is a compact, orientable surface of genus γ then :

$$\lambda_k(H) - \lambda_1(H) \leq C \left(\frac{\|\psi\|_\infty}{\|\psi\|_2^g} \right)^2 (\gamma + 1)k$$

where C is an absolute constant.

Proof. As $H = D^*D + T$, the quadratic form associated to H is given by:

$$\mathcal{Q}(\psi) = \int_M \langle H\psi, \psi \rangle dv_g = \int_M \left(|D\psi|^2 + \langle T\psi, \psi \rangle \right) dv_g,$$

where ψ denotes a generic smooth section. If u is any Lipschitz function on M , then an integration by parts gives (see [6, Lemma 8])

$$\mathcal{Q}(u\psi) = \int_M \left(u^2 \langle H\psi, \psi \rangle + |\nabla u|^2 |\psi|^2 \right) dv_g.$$

Now assume that ψ is a first eigensection: $H\psi = \lambda_1(H)\psi$. Then we obtain:

$$\frac{\mathcal{Q}(u\psi)}{\int_M u^2 |\psi|^2 dv_g} = \lambda_1(H) + \frac{\int_M |\nabla u|^2 |\psi|^2 dv_g}{\int_M u^2 |\psi|^2 dv_g}$$

for all Lipschitz functions u . Let $\sigma = |\psi|^2$ and $\nu = |\psi|^2 dv_g$. Restricting the test-sections to sections of type $u\psi$, where u is Lipschitz (hence in $H^1(M)$) and ψ is a fixed first eigensection, an obvious application of the min-max principle gives:

$$\lambda_k(H) \leq \lambda_1(H) + \mu_k(\sigma, \nu).$$

The remaining part of the theorem is an immediate consequence of the last inequality and Theorem 1.1. \square

Corollary 4.1. *Assume that a Laplace type operator H acting on sections of a Riemannian vector bundle E over (M, g) , admits a first eigensection of constant length. Then, for all $k \geq 1$*

$$\lambda_k(H) - \lambda_1(H) \leq \lambda_k(\Delta_g).$$

Indeed, when σ is constant, $\mu_k(\sigma, \sigma dv_g)$ is nothing but the k -th eigenvalue of the Laplacian Δ_g acting on functions. In the particular case where $H^{(p)}$ is the Hodge Laplacian acting on p -forms, Corollary 4.1 says that the existence of a nonzero harmonic p -form of constant length on M leads to

$$\lambda_k(H^{(p)}) \leq \lambda_k(\Delta_g)$$

for every positive integer k , which extends the result of Takahashi [42].

5. LOWER BOUNDS FOR EIGENVALUES ON WEIGHTED EUCLIDEAN DOMAINS AND MANIFOLDS OF REVOLUTION

The scope of this section is to show that our main upper bound is asymptotically sharp for some special weighted manifolds, namely convex Euclidean domains and revolution manifolds endowed with a Gaussian density. By a Gaussian density we mean a function of type

$$\sigma_j(x) = e^{-jd(x, x_0)^2}$$

where x_0 is a fixed point and j is a positive integer (a slight modification is needed for closed revolution manifolds). We will give a lower bound of $\lambda_2(L_{\sigma_j})$ and verify that in all these cases both $\lambda_2(L_{\sigma_j})$ and our main upper bound grow to infinity linearly in j as $j \rightarrow \infty$.

In what follows, we make use of the Reilly formula, recently extended to weighted manifolds (see [33]). Here M is a compact Riemannian manifold

of dimension n with smooth boundary ∂M ; we let $\sigma = e^{-f}$ be a positive smooth density and $u \in C^\infty(M)$. Then we have:

$$\begin{aligned} \int_M (L_\sigma u)^2 \sigma &= \int_M |\nabla^2 u|^2 \sigma + \text{Ric}(\nabla u, \nabla u) \sigma + \nabla^2 f(\nabla u, \nabla u) \sigma \\ &\quad + \int_{\partial M} 2 \frac{\partial u}{\partial N} L_\sigma^{\partial M} u \cdot \sigma + B(\nabla^{\partial M} u, \nabla^{\partial M} u) \sigma \\ &\quad + \int_{\partial M} \left((n-1)H + \frac{\partial f}{\partial N} \right) \left(\frac{\partial u}{\partial N} \right)^2 \sigma \end{aligned} \quad (13)$$

where N is the inner unit normal, B the second fundamental form of ∂M with respect to N and $(n-1)H = \text{tr} B$ is the mean curvature. By $L_\sigma^{\partial M}$ we denote the induced operator on ∂M , naturally defined as

$$L_\sigma^{\partial M} u = \Delta^{\partial M} u + \langle \nabla^{\partial M} f, \nabla^{\partial M} u \rangle.$$

When not explicitly indicated, integration is taken with respect to the canonical Riemannian measure.

5.1. Convex Euclidean domains. Here is the main statement of this section.

Theorem 5.1. *Let Ω be a convex domain of \mathbb{R}^n containing the origin, endowed with the Gaussian density $\sigma_j(x) = e^{-j|x|^2}$. There exists a constant $K_{n,R}$ depending only on n and $R = \text{dist}(O, \partial\Omega)$ such that, for all $j \geq K_{n,R}$,*

$$\frac{\|\sigma_j\|_{\frac{n}{n-2}}}{\|\sigma_j\|_1} \leq \lambda_2(L_{\sigma_j}) \leq B_n \frac{\|\sigma_j\|_{\frac{n}{n-2}}}{\|\sigma_j\|_1} \quad (14)$$

where B_n is a constant that depends only on n . Moreover, $\lambda_2(L_{\sigma_j})$ tends to infinity with a linear growth as $j \rightarrow \infty$.

The proof follows from Theorem 1.1 and the following estimates.

Proposition 5.1. *Let Ω be a convex domain in \mathbb{R}^n . Then, for all $j > 0$,*

$$\lambda_2(L_{\sigma_j}) \geq 2j.$$

Proof. We apply the Reilly formula (13) to an eigenfunction u associated to $\lambda = \lambda_2(L_{\sigma_j})$. By definition, $f = j|x|^2$ so that $\nabla^2 f = 2jI$. As $\frac{\partial u}{\partial N} = 0$ (Neumann condition) and $B \geq 0$ by the convexity assumption, we arrive at

$$\lambda^2 \int_\Omega u^2 \sigma \geq 2j \int_\Omega |\nabla u|^2 \sigma = 2j\lambda \int_\Omega u^2 \sigma,$$

and the inequality follows. \square

Lemma 5.1. *Let $j \in \mathbb{N}^*$.*

- a) *For all $p > 1$ one has $\|\sigma_j\|_p \leq \frac{C_{n,p}}{j^{\frac{n}{2p}}}$. In particular $\|\sigma_j\|_{\frac{n}{n-2}} \leq \frac{C_n}{j^{\frac{n-2}{2}}}$.*
- b) *There exists a constant $K_{n,R}$ depending only on n and R such that, if $j \geq K_{n,R}$, then $\|\sigma_j\|_1 \geq \frac{C_n}{2j^{\frac{n}{2}}}$.*

The constants $C_n, C_{n,p}$ will be given explicitly in the proof. From the above facts we obtain for $j \geq K_{n,R}$,

$$\frac{\|\sigma_j\|_{\frac{n}{n-2}}}{\|\sigma_j\|_1} \leq 2j \leq \lambda_2(L_{\sigma_j}).$$

Proof of Lemma 5.1. We use the well-known formula

$$\int_0^\infty r^{n-1} e^{-jr^2} dr = \frac{\Gamma(\frac{n}{2})}{2j^{\frac{n}{2}}},$$

where Γ is the Gamma function. Then

$$\int_{\mathbb{R}^n} e^{-j|x|^2} dx = |\mathbb{S}^{n-1}| \int_0^\infty r^{n-1} e^{-jr^2} dr = \frac{C_n}{j^{\frac{n}{2}}}.$$

with $C_n = \frac{1}{2} |\mathbb{S}^{n-1}| \Gamma(\frac{n}{2})$. This implies that

$$\|\sigma_j\|_p = \left(\int_{\Omega} e^{-jp|x|^2} dx \right)^{\frac{1}{p}} \leq \left(\int_{\mathbb{R}^n} e^{-jp|x|^2} dx \right)^{\frac{1}{p}} = \frac{C_{n,p}}{j^{\frac{n}{2p}}}$$

with $C_{n,p} = \frac{(C_n)^{\frac{1}{p}}}{p^{\frac{n}{2p}}}$, which proves a). Now, one has

$$\|\sigma_j\|_1 = \int_{\Omega} e^{-j|x|^2} dx = \int_{\mathbb{R}^n} e^{-j|x|^2} dx - \int_{\Omega^c} e^{-j|x|^2} dx = \frac{C_n}{j^{\frac{n}{2}}} - \int_{\Omega^c} e^{-j|x|^2} dx.$$

On the complement of Ω one has $|x| \geq R$ by the definition of R . Then,

$$\int_{\Omega^c} e^{-j|x|^2} dx = \int_{\Omega^c} e^{-\frac{j}{2}|x|^2} e^{-\frac{j}{2}|x|^2} dx \leq e^{-\frac{jR^2}{2}} \int_{\mathbb{R}^n} e^{-\frac{|x|^2}{2}} dx = D_n e^{-\frac{jR^2}{2}}.$$

As R is fixed and $j^{\frac{n}{2}} e^{-\frac{jR^2}{2}}$ tends to zero when j tends to infinity, one sees immediately that there exists $K_{n,R}$ such that, for $j \geq K_{n,R}$,

$$\int_{\Omega^c} e^{-j|x|^2} dx \leq D_n e^{-\frac{jR^2}{2}} \leq \frac{C_n}{2j^{\frac{n}{2}}}.$$

In that range of j one indeed has

$$\|\sigma_j\|_1 \geq \frac{C_n}{2j^{\frac{n}{2}}}.$$

□

5.2. Revolution manifolds with boundary. A *revolution manifold with boundary* of dimension n is a Riemannian manifold (M, g) with a distinguished point N such that $(M \setminus \{N\}, g)$ is isometric to $(0, R] \times \mathbb{S}^{n-1}$ endowed with the metric

$$g = dr^2 + \theta^2(r) g_{\mathbb{S}^{n-1}},$$

$g_{\mathbb{S}^{n-1}}$ denoting the standard metric on the sphere. Here $\theta(r)$ is a smooth function on $[0, R]$ which is positive on $(0, R]$ (note that we assume in particular $\theta(R) > 0$) and is such that:

$$\theta(0) = \theta''(0) = 0, \theta'(0) = 1.$$

We notice that, as $R < +\infty$, M is compact, connected and has a smooth boundary $\partial M = \{R\} \times \mathbb{S}^{n-1}$ isometric to the $(n-1)$ -dimensional sphere

of radius $\theta(R)$. If we make the stronger assumption that θ has vanishing even derivatives at zero then the metric is C^∞ -smooth everywhere. Given a Gaussian radial density of the form $\sigma_j(r) = e^{-jr^2}$ centered at the pole N , we wish to study the first non-zero eigenvalue $\lambda_2(L_{\sigma_j})$ of L_{σ_j} , with Neumann boundary conditions.

Theorem 5.2. *Let (M, g) be a revolution manifold with boundary endowed with the Gaussian density $\sigma_j(r) = e^{-jr^2}$, $j \geq 1$. Then there is a (possibly negative) constant C , not depending on j , such that*

$$\lambda_2(L_{\sigma_j}) \geq 2j + C.$$

Moreover, if (M, g) has non-negative Ricci curvature, then $\lambda_2(L_{\sigma_j}) \geq 2j$.

We start the proof by making general considerations which are valid for any radial density $\sigma(r) = e^{-f(r)}$ (Lemma 5.2 and 5.3). As for the usual Laplacian, we can separate variables and prove that there is an orthonormal basis of $L^2(\sigma)$ made of eigenfunctions of type

$$u(r, x) = \phi(r)\xi(x) \quad (15)$$

where ϕ is a smooth function of $r \in [0, R]$ and $\xi(x)$ is an eigenfunction of the Laplacian on \mathbb{S}^{n-1} . Listing the eigenvalues of \mathbb{S}^{n-1} as $\{\mu_k\}$, where $k = 1, 2, \dots$, and computing the Laplacian, one arrives at

$$\Delta u(r, x) = \left(-\phi''(r) - (n-1)\frac{\theta'(r)}{\theta(r)}\phi'(r) + \frac{\mu_k}{\theta(r)^2}\phi(r) \right) \xi(x).$$

As $f = f(r)$ is radial one has $\langle \nabla f, \nabla u \rangle(r, x) = f'(r)\phi'(r)\xi(x)$ so that

$$L_\sigma u(r, x) = \left(-\phi''(r) - (n-1)\frac{\theta'(r)}{\theta(r)}\phi'(r) + f'(r)\phi'(r) + \frac{\mu_k}{\theta(r)^2}\phi(r) \right) \xi(x) \quad (16)$$

Let us now focus on the first positive eigenvalue $\lambda_2(L_\sigma)$ with associated eigenfunction u of the form (15). There are only two cases to examine:

- either $\mu_k = \mu_1 = 0$, so that $\xi(x)$ is constant and the eigenfunction u is *radial*,
- or $\mu_k = \mu_2 = n-1$, the first positive eigenvalue of \mathbb{S}^{n-1} .

In fact, higher eigenvalues of \mathbb{S}^{n-1} do not occur, otherwise u would have too many nodal domains. We summarize this alternative in the following lemma.

Lemma 5.2. *Let $M = (0, R] \times \mathbb{S}^{n-1}$ be a manifold of revolution as above, endowed with a radial density $\sigma(r) = e^{-f(r)}$. Then, either L_σ admits a (Neumann) radial eigenfunction associated to $\lambda_2(L_\sigma)$, or $\lambda_2(L_\sigma)$ is the first positive eigenvalue of the problem*

$$\begin{cases} \phi'' + \left((n-1)\frac{\theta'}{\theta} - f' \right) \phi' + \left(\lambda - \frac{n-1}{\theta^2} \right) \phi = 0 \\ \phi(0) = \phi'(R) = 0 \end{cases}$$

Note that the condition $\phi'(R) = 0$ follows from the Neumann boundary condition, while the condition $\phi(0) = 0$ is imposed to insure that the eigenfunction is continuous at the pole N .

We then prove the following lower bound for the "radial spectrum".

Lemma 5.3. *In the hypothesis of Lemma 5.2, assume that L_σ admits a a (Neumann) radial eigenfunction associated to the eigenvalue λ . Then*

$$\lambda \geq \inf_{(0,R)} \left\{ (n-1) \left(\frac{\theta'}{\theta} \right)^2 + \text{Ric}_0 + f'' \right\},$$

where Ric_0 is a lower bound of the Ricci curvature of M .

Proof. Let $u = u(r)$ be a radial eigenfunction associated to λ . We apply the Reilly formula (13) to obtain:

$$\lambda \int_M u^2 \sigma = \int_M |\nabla^2 u|^2 \sigma + \text{Ric}(\nabla u, \nabla u) \sigma + \nabla^2 f(\nabla u, \nabla u) \sigma.$$

In fact, the boundary terms vanish because on ∂M one has $\frac{\partial u}{\partial N} = 0$ and, as u is radial (hence constant on ∂M), one also has $\nabla^{\partial M} u = 0$. We now wish to bound from below the terms involving the Hessians. For that we need to use a suitable orthonormal frame. So, fix a point $p = (r, x)$ and consider a local frame $(\bar{e}_1, \dots, \bar{e}_{n-1})$ around x which is orthonormal for the canonical metric of \mathbb{S}^{n-1} . We can assume that this frame is geodesic at x . Taking $e_i = \frac{1}{\theta} \bar{e}_i$ it is clear that $(e_1, \dots, e_{n-1}, \frac{\partial}{\partial r})$ is a local orthonormal frame on (M, g) . If ∇ denotes (as usual) the Levi-Civita connection of (M, g) one sees easily that at p ,

$$\nabla_{e_i} e_j = -\delta_{ij} \frac{\theta'}{\theta} \frac{\partial}{\partial r}, \quad \nabla_{e_i} \frac{\partial}{\partial r} = \frac{\theta'}{\theta} e_i, \quad \nabla_{\frac{\partial}{\partial r}} e_i = \nabla_{\frac{\partial}{\partial r}} \frac{\partial}{\partial r} = 0.$$

Since $\nabla u = u' \frac{\partial}{\partial r}$ we have $\langle \nabla u, e_i \rangle = 0$ for all i . Then

$$\nabla^2 u(e_i, e_j) = \langle \nabla_{e_i} \nabla u, e_j \rangle = e_i \cdot \langle \nabla u, e_j \rangle - \langle \nabla u, \nabla_{e_i} e_j \rangle = \delta_{ij} \frac{\theta'}{\theta} u'.$$

Similarly, one shows that $\nabla^2 u(e_i, \frac{\partial}{\partial r}) = 0$ and $\nabla^2 u(\frac{\partial}{\partial r}, \frac{\partial}{\partial r}) = u''$. It follows that the matrix of $\nabla^2 u$ in the given basis is diagonal, that is

$$\nabla^2 u = \text{diag} \left(\frac{\theta'}{\theta} u', \dots, \frac{\theta'}{\theta} u', u'' \right)$$

which in turn implies that

$$|\nabla^2 u|^2 \geq (n-1) \left(\frac{\theta'}{\theta} \right)^2 u'^2 = (n-1) \left(\frac{\theta'}{\theta} \right)^2 |\nabla u|^2.$$

On the other hand

$$\nabla^2 f(\nabla u, \nabla u) = u'^2 \nabla^2 f \left(\frac{\partial}{\partial r}, \frac{\partial}{\partial r} \right) = f'' |\nabla u|^2.$$

Substituting in the Reilly formula above, we obtain

$$\lambda^2 \int_M u^2 \sigma \geq \int_M \left((n-1) \left(\frac{\theta'}{\theta} \right)^2 + \text{Ric}_0 + f'' \right) |\nabla u|^2 \sigma$$

so that, if

$$C = \inf_{(0,R)} \left\{ n \left(\frac{\theta'}{\theta} \right)^2 + \text{Ric}_0 + f'' \right\}$$

then

$$\lambda^2 \int_M u^2 \sigma \geq C \int_M |\nabla u|^2 \sigma = C \lambda \int_M u^2 \sigma$$

which implies $\lambda \geq C$ as asserted. \square

We are now ready to prove Theorem 5.2. Let $\sigma_j = e^{-jr^2}$ so that $f''(r) = 2j$. If $\lambda_2(L_{\sigma_j})$ is associated to a radial eigenfunction, Lemma 5.3 gives

$$\lambda_2(L_{\sigma_j}) \geq 2j + C_1,$$

with C_1 independent of j . It is also clear that if M has non-negative Ricci curvature then $\lambda_2(L_{\sigma_j}) \geq 2j$.

Taking into account Lemma 5.2, Theorem 5.2 will now follow from

Lemma 5.4. *Let λ_2 be the first positive eigenvalue of the problem*

$$\begin{cases} \phi'' + \left((n-1)\frac{\theta'}{\theta} - 2jr\right)\phi' + \left(\lambda - \frac{n-1}{\theta^2}\right)\phi = 0 \\ \phi(0) = \phi'(R) = 0 \end{cases}$$

Then

$$\lambda_2 \geq 2j + C_2$$

where $C_2 = (n-1) \inf \left\{ \frac{r - \theta'\theta}{r\theta^2} : r \in (0, R) \right\}$. Moreover, if M has non-negative Ricci curvature, then $C_2 \geq 0$.

We observe that C_2 is always finite because since $r \rightarrow \frac{r - \theta'\theta}{r\theta^2}$ approaches $-\frac{2}{3}\theta'''(0)$ as $r \rightarrow 0$, and then is bounded on $[0, R]$.

Proof. Set $\phi(r) = ry(r)$ so that $\phi' = y + ry'$ and $\phi'' = 2y' + ry''$. Substituting in the equation we obtain

$$y'' + \left(\frac{2}{r} + (n-1)\frac{\theta'}{\theta} - 2jr\right)y' + \left(\lambda - 2j + (n-1)\frac{\theta'\theta - r}{r\theta^2}\right)y = 0.$$

If $\beta = r^2\theta^{n-1}e^{-jr^2}$, then $\frac{2}{r} + (n-1)\frac{\theta'}{\theta} - 2jr = \frac{\beta'}{\beta}$ and the equation takes the form

$$(\beta y')' + \left(\lambda - 2j + (n-1)\frac{\theta'\theta - r}{r\theta^2}\right)\beta y = 0.$$

Multiplying by y and integrating on $(0, R)$ we end-up with

$$(\lambda - 2j) \int_0^R \beta y^2 dr = - \int_0^R (\beta y')' y dr + \int_0^R (n-1) \frac{r - \theta'\theta}{r\theta^2} \beta y^2 dr.$$

We observe that $\beta(0) = 0$ and, as $\phi'(R) = 0$, we have $y(R) = -Ry'(R)$. Then, integrating by parts:

$$\begin{aligned} - \int_0^R (\beta y')' y dr &= -\beta(R)y'(R)y(R) + \int_0^R \beta y'^2 dr \\ &= \beta(R)y'(R)^2 R + \int_0^R \beta y'^2 dr \geq 0. \end{aligned}$$

From the definition of the constant C_2 we obtain

$$(\lambda - 2j) \int_0^R \beta y^2 dr \geq C_2 \int_0^R \beta y^2 dr$$

which gives the assertion.

Finally, it is well-known that $\text{Ric}(\frac{\partial}{\partial r}, \frac{\partial}{\partial r}) = -(n-1)\frac{\theta''}{\theta}$. If M has non-negative Ricci curvature then $\theta'' \leq 0$; in turn we have $\theta' \leq 1$ and $\theta \leq r$, which implies that $r - \theta'\theta \geq 0$ and, then, $C_2 \geq 0$. \square

5.3. Closed revolution manifolds. A *closed revolution manifold* of dimension n is a compact manifold (M, g) without boundary, having two distinguished points N, S such that $(M \setminus \{N, S\}, g)$ is isometric to $(0, R) \times \mathbb{S}^{n-1}$ endowed with the metric $dr^2 + \theta^2(r)g_{\mathbb{S}^{n-1}}$, where $\theta : [0, R] \rightarrow \mathbb{R}$ is smooth and

$$\theta(0) = \theta(R) = 0, \quad \theta'(0) = -\theta'(R) = 1, \quad \theta''(0) = \theta''(R) = 0.$$

Under these assumptions, the metric g is C^2 . To have a C^∞ metric it is enough to assume that $\theta^{(2i)}(0) = \theta^{(2i)}(R) = 0$ for all $i = 0, 1, 2, \dots$.

Our aim is to construct a sequence of radial densities $\{\sigma_j\}$ on M such that $\lambda_2(L_{\sigma_j})$ grows linearly with j . These densities will be Gaussian functions centered at the pole N of M , suitably smoothened near the pole S so that the resulting function is globally C^1 (this is enough for our purpose). Thus, let us define

$$\sigma_j(r) = e^{-f_j(r)} \tag{17}$$

where $f_j(r) = jh_j^2(r)$ and

$$h_j(r) = \begin{cases} r & \text{if } 0 \leq r \leq r_j \doteq R - \frac{1}{\alpha j} \\ r - \frac{\alpha j}{2}(r - r_j)^2 & \text{if } r_j \leq r \leq R \end{cases} \tag{18}$$

Here $\alpha \geq 1$ is a fixed number, which is large enough so that

$$\frac{(n-1)\alpha^2}{16} - 2\alpha R \geq 2.$$

The function h_j is of class C^1 on $[0, R]$. Since $\sigma_j'(0) = \sigma_j'(R) = 0$, we see that σ_j is a C^1 function on M .

Theorem 5.3. *Let (M, g) be a closed revolution manifold endowed with the density σ_j defined in (17) and (18). Then there exist an integer j_0 and a constant C , not depending on j , such that, for all $j \geq j_0$, one has:*

$$\lambda_2(L_{\sigma_j}) \geq 2j + C.$$

We give the proof of Theorem 5.3 in the next subsection. We just want to mention here the analogue of Theorem 5.1.

Theorem 5.4. *Let (M, g) be a revolution manifold with boundary (resp. a closed revolution manifold) endowed with the density σ_j as in Theorem 5.2 (resp. Theorem 5.3). Then, for j sufficiently large,*

$$A \frac{\|\sigma_j\|_{\frac{n}{n-2}}}{\|\sigma_j\|_1} \leq \lambda_2(L_{\sigma_j}) \leq B \frac{\|\sigma_j\|_{\frac{n}{n-2}}}{\|\sigma_j\|_1}$$

where A and B are positive constants depending on M , but not on j . Moreover, $\lambda_2(L_{\sigma_j})$ tends to infinity with a linear growth as $j \rightarrow \infty$.

The proof of Theorem 5.4 is quite similar to that of the corresponding statement in Theorem 5.1, and we only sketch it. Assume that M has a boundary and $n \geq 3$. We start from:

$$\|\sigma_j\|_p^p = \int_M \sigma_j^p(r) dv_g = |\mathbb{S}^{n-1}| \int_0^R \theta^{n-1}(r) \sigma_j^p(r) dr.$$

Now $|\theta(r) - r| \leq cr^3$ and $\theta^{n-1}(r) \sim r^{n-1}$ as $r \rightarrow 0$; since, for $p = \frac{n}{n-2}$ and j large:

$$\frac{\left(\int_0^R r^{n-1} e^{-jpr^2} dr \right)^{\frac{1}{p}}}{\int_0^R r^{n-1} e^{-jr^2} dr} \leq C' j,$$

we get in turn

$$\frac{\|\sigma_j\|_{\frac{n}{n-2}}}{\|\sigma_j\|_1} \leq C'' j.$$

The assertion follows from the last inequality and the estimate of $\lambda_2(L_{\sigma_j})$ we have obtained in Theorem 5.2. If M is closed the only change is in the definition of σ_j . However, this change occurs far from the pole and thus contributes with exponentially decreasing terms, which, modulo a change in the constants, do not show up in the final estimate. We omit further details.

5.4. Proof of Theorem 5.3. We start the proof by observing, as in the previous section, that there is an eigenfunction associated to $\lambda_2(L_{\sigma_j})$ of type $u(r, x) = \phi(r)\xi(x)$ with ϕ satisfying a suitable Sturm-Liouville problem on the interval $[0, R]$ and ξ being an eigenfunction of the Laplacian on \mathbb{S}^{n-1} . For a closed manifold, Lemma 5.2 takes the following form.

Lemma 5.5. *Let (M, g) be a closed revolution manifold endowed with a radial density $\sigma(r) = e^{-f(r)}$. Then, either L_σ admits a radial eigenfunction associated to $\lambda_2(L_\sigma)$, or $\lambda_2(L_\sigma)$ is the first positive eigenvalue of the following Sturm-Liouville problem on $[0, R]$:*

$$\begin{cases} \phi'' + \left((n-1) \frac{\theta'}{\theta} - f' \right) \phi' + \left(\lambda - \frac{n-1}{\theta^2} \right) \phi = 0 \\ \phi(0) = \phi(R) = 0 \end{cases}$$

Note the boundary conditions on ϕ , which are imposed so that the corresponding eigenfunction u is continuous at both poles N and S .

By definition, $R - r_j = \frac{1}{\alpha_j}$ and, as $j \rightarrow \infty$ we have $r_j \rightarrow R$. Since θ' is continuous and $\theta'(R) = -1$, there exists an integer j_1 such that, for all $j \geq j_1$,

$$-2 \leq \theta'(r) \leq -\frac{1}{2}$$

for all $r \in [r_j, R]$. Consequently, on that interval we also have

$$\frac{1}{2}(R - r) \leq \theta(r) \leq 2(R - r) \quad \text{and} \quad \left| \frac{\theta'(r)}{\theta(r)} \right| \geq \frac{1}{4(R - r)}. \quad (19)$$

According to Lemma 5.5 there are two cases to discuss.

First case: there is a radial eigenfunction associated to $\lambda = \lambda_2(L_{\sigma_j})$.

Then, we apply Lemma 5.3 (which holds without change) and for the lower bound it suffices to verify the inequality:

$$(n-1)\left(\frac{\theta'(r)}{\theta(r)}\right)^2 + f_j''(r) \geq 2j \quad (20)$$

for all $r \in (0, R)$. Indeed, one has $f_j'' = 2j(h_j'^2 + h_j h_j'')$. Thus, on the interval $(0, r_j)$ one gets $f_j'' = 2j$ and (20) is immediate. On the interval (r_j, R) one has $h_j \leq r \leq R$ and $h_j'' = -\alpha j$. Then:

$$f_j'' \geq 2j h_j h_j'' \geq -2\alpha j^2 R.$$

On the other hand, as $R - r \leq R - r_j = \frac{1}{\alpha j}$, we see from (19):

$$\left(\frac{\theta'(r)}{\theta(r)}\right)^2 \geq \frac{1}{16(R-r)^2} \geq \frac{\alpha^2 j^2}{16}.$$

Recalling the definition of α we see that, for $j \geq j_1$:

$$(n-1)\left(\frac{\theta'(r)}{\theta(r)}\right)^2 + f_j''(r) \geq \left((n-1)\frac{\alpha^2}{16} - 2\alpha R\right)j^2 \geq 2j^2 \geq 2j,$$

and the assertion follows.

Second case: $\lambda_2(L_{\sigma_j})$ is the first positive eigenvalue of the problem:

$$\begin{cases} \phi'' + \left((n-1)\frac{\theta'}{\theta} - f_j'\right)\phi' + \left(\lambda - \frac{n-1}{\theta^2}\right)\phi = 0 \\ \phi(0) = \phi(R) = 0 \end{cases}$$

First observe that there exists $\bar{R} \in (0, R)$ such that $\phi'(\bar{R}) = 0$ (note that \bar{R} depends on j). It follows that u is a Neumann eigenfunction associated to $\lambda_2(L_{\sigma_j})$ for both of the following domains:

$$\Omega_1 = \{(r, x) \in M : r \leq \bar{R}\}, \quad \Omega_2 = \{(r, x) \in M : r \geq \bar{R}\}.$$

We first assume that $\bar{R} < r_j$ and focus our attention on Ω_1 . As $f_j' = 2jr$ on $[0, \bar{R}]$, we see that $\lambda_2(L_{\sigma_j})$ is bounded below by the first positive eigenvalue of the problem

$$\begin{cases} \phi'' + \left((n-1)\frac{\theta'}{\theta} - 2jr\right)\phi' + \left(\lambda - \frac{n-1}{\theta^2}\right)\phi = 0 \\ \phi(0) = \phi'(\bar{R}) = 0. \end{cases}$$

By Lemma 5.4 we have

$$\lambda_2(L_{\sigma_j}) \geq 2j + \tilde{C}_2,$$

where $\tilde{C}_2 = (n-1) \inf_{0 < r > \bar{R}} \left\{ \frac{r - \theta'\theta}{r\theta^2} \right\} \geq (n-1) \inf_{0 < r > R} \left\{ \frac{r - \theta'\theta}{r\theta^2} \right\}$, a constant which does not depend on j , and we are done.

Finally, it remains to examine the case where $\bar{R} \geq r_j$. In this case, we view u as a Neumann eigenfunction on the domain Ω_2 . Let us briefly sketch the argument. As $j \rightarrow \infty$, Ω_2 is quasi-isometric to a Euclidean ball of small radius (of the order of $1/j$) and has a density with uniformly controlled

variation. Therefore, its first positive eigenvalue must be large (of the order of j^2). Let us clarify the details. We know that

$$\lambda_2(L_{\sigma_j}) = \frac{\int_{\Omega_2} |\nabla u|^2 \sigma_j dv_g}{\int_{\Omega_2} u^2 \sigma_j dv_g}.$$

Since $h_j(r)$ is increasing in $r \in (r_j, R)$, one has $r_j = h_j(r_j) \leq h_j(r) \leq h_j(R) \leq R$. Thus, for all $r \in (\bar{R}, R)$, one has (recall that $r_j = R - \frac{1}{\alpha j}$)

$$e^{-jR^2} \leq \sigma_j(r) \leq e^{-j(R - \frac{1}{\alpha j})^2}.$$

Consequently,

$$\lambda_2(L_{\sigma_j}) \geq \frac{\int_{\Omega_2} |\nabla u|^2 dv_g}{\int_{\Omega_2} u^2 dv_g} e^{-\frac{2R}{\alpha}}. \quad (21)$$

As $u(r, x) = \phi(r)\xi(x)$ and $\int_{\mathbb{S}^{n-1}} \xi = 0$, we see that $\int_{\Omega_2} u dv_g = 0$. Hence, by the min-max principle:

$$\frac{\int_{\Omega_2} |\nabla u|^2 dv_g}{\int_{\Omega_2} u^2 dv_g} \geq \mu_2(\Omega_2, g) \quad (22)$$

where $\mu_2(\Omega_2, g)$ denotes the first positive Neumann eigenvalue of the Laplacian on the domain (Ω_2, g) . On the other hand, the first inequality in (19) shows that the metric g is quasi-isometric, on Ω_2 , to the standard Euclidean metric $g_{euc} = dr^2 + r^2 g_{\mathbb{S}^{n-1}}$, with quasi-isometry ratio bounded by 4. Thus, (Ω_2, g) is quasi-isometric to the Euclidean ball (Ω_2, g_{euc}) of radius $R - \bar{R}$. Therefore:

$$\mu_2(\Omega_2, g) \geq 4^{-(n+2)} \mu_2(\Omega_2, g_{euc}) \geq \frac{4^{-(n+2)} \mu(n+1)}{(R - \bar{R})^2} \quad (23)$$

where $\mu(n+1)$ is the first positive Neumann eigenvalue of the unit Euclidean ball. Since $R - \bar{R} \leq R - r_j = \frac{1}{\alpha j}$, we conclude from (21), (22), (23) that:

$$\lambda_2(L_{\sigma_j}) \geq C_3 j^2$$

where C_3 is a constant depending only on n, α and R . By taking j larger than a suitable integer j_2 we see that $\lambda_2(L_{\sigma_j}) \geq 2j$ as asserted. The conclusion is that, if $j \geq j_0 \doteq \max\{j_1, j_2\}$, then the inequality of Theorem 5.3 is verified. This ends the proof.

REFERENCES

- [1] Ben Andrews and Lei Ni. Eigenvalue comparison on Bakry-Emery manifolds. *Comm. Partial Differential Equations*, 37(11):2081–2092, 2012.
- [2] D. Bakry and Michel Émery. Diffusions hypercontractives. In *Séminaire de probabilités, XIX, 1983/84*, volume 1123 of *Lecture Notes in Math.*, pages 177–206. Springer, Berlin, 1985.
- [3] Pierre H. Bérard. From vanishing theorems to estimating theorems: the Bochner technique revisited. *Bull. Amer. Math. Soc. (N.S.)*, 19(2):371–406, 1988.
- [4] B. Colbois and J. Dodziuk. Riemannian metrics with large λ_1 . *Proc. Amer. Math. Soc.*, 122(3):905–906, 1994.
- [5] Bruno Colbois and Ahmad El Soufi. Eigenvalues of the Laplacian acting on p -forms and metric conformal deformations. *Proc. Amer. Math. Soc.*, 134(3):715–721 (electronic), 2006.
- [6] Bruno Colbois and Alessandro Savo. Large eigenvalues and concentration. *Pacific J. Math.*, 249(2):271–290, 2011.

- [7] Bruno Colbois and Alessandro Savo. Involutive isometries, eigenvalue bounds and a spectral property of Clifford tori. *Indiana Univ. Math. J.*, 61(1):337–357, 2012.
- [8] A. El Soufi and S. Ilias. Immersions minimales, première valeur propre du laplacien et volume conforme. *Math. Ann.*, 275(2):257–267, 1986.
- [9] Ahmad El Soufi, Hector Giacomini, and Mustapha Jazar. A unique extremal metric for the least eigenvalue of the Laplacian on the Klein bottle. *Duke Math. J.*, 135(1):181–202, 2006.
- [10] Ahmad El Soufi, Saïd Ilias, and Antonio Ros. Sur la première valeur propre des tores. In *Séminaire de Théorie Spectrale et Géométrie, No. 15, Année 1996–1997*, volume 15 of *Sémin. Théor. Spectr. Géom.*, pages 17–23. Univ. Grenoble I, Saint-Martin-d’Hères, 1997.
- [11] Kei Funano and Takashi Shioya. Concentration, Ricci Curvature, and Eigenvalues of Laplacian. *Geom. Funct. Anal.*, 23(3):888–936, 2013.
- [12] Akito Futaki, Haizhong Li, and Xiang-Dong Li. On the first eigenvalue of the Witten-Laplacian and the diameter of compact shrinking ricci solitons. *arXiv:1111.6364*, 2011.
- [13] Akito Futaki and Yuji Sano. Lower diameter bounds for compact shrinking Ricci solitons. *Asian J. Math.*, 2010.
- [14] Peter B. Gilkey, John V. Leahy, and Jeonghyeong Park. *Spectral geometry, Riemannian submersions, and the Gromov-Lawson conjecture*. Studies in Advanced Mathematics. Chapman & Hall/CRC, Boca Raton, FL, 1999.
- [15] Alexandre Girouard, Nikolai Nadirashvili, and Iosif Polterovich. Maximization of the second positive Neumann eigenvalue for planar domains. *J. Differential Geom.*, 83(3):637–661, 2009.
- [16] A. Grigor’yan and S.-T. Yau. Decomposition of a metric space by capacitors. In *Differential equations: La Pietra 1996 (Florence)*, volume 65 of *Proc. Sympos. Pure Math.*, pages 39–75. Amer. Math. Soc., Providence, RI, 1999.
- [17] Alexander Grigor’yan. Heat kernels on weighted manifolds and applications. In *The ubiquitous heat kernel*, volume 398 of *Contemp. Math.*, pages 93–191. Amer. Math. Soc., Providence, RI, 2006.
- [18] Alexander Grigor’yan, Yuri Netrusov, and Shing-Tung Yau. Eigenvalues of elliptic operators and geometric applications. In *Surveys in differential geometry. Vol. IX*, *Surv. Differ. Geom.*, IX, pages 147–217. Int. Press, Somerville, MA, 2004.
- [19] M. Gromov and V. D. Milman. A topological application of the isoperimetric inequality. *Amer. J. Math.*, 105(4):843–854, 1983.
- [20] Asma Hassannezhad. Conformal upper bounds for the eigenvalues of the Laplacian and Steklov problem. *J. Funct. Anal.*, 261(12):3419–3436, 2011.
- [21] Asma Hassannezhad. Eigenvalues of perturbed Laplace operators on compact manifolds. *Pacific J. Math.*, 2012.
- [22] B. Helffer and J. Sjöstrand. Puits multiples en mécanique semi-classique. IV. Étude du complexe de Witten. *Comm. Partial Differential Equations*, 10(3):245–340, 1985.
- [23] Bernard Helffer, Markus Klein, and Francis Nier. Quantitative analysis of metastability in reversible diffusion processes via a Witten complex approach. *Mat. Contemp.*, 26:41–85, 2004.
- [24] Joseph Hersch. Quatre propriétés isopérimétriques de membranes sphériques homogènes. *C. R. Acad. Sci. Paris Sér. A-B*, 270:A1645–A1648, 1970.
- [25] Dmitry Jakobson, Nikolai Nadirashvili, and Iosif Polterovich. Extremal metric for the first eigenvalue on a Klein bottle. *Canad. J. Math.*, 58(2):381–400, 2006.
- [26] Nicholas Korevaar. Upper bounds for eigenvalues of conformal metrics. *J. Differential Geom.*, 37(1):73–93, 1993.
- [27] Peter Li and Shing Tung Yau. A new conformal invariant and its applications to the Willmore conjecture and the first eigenvalue of compact surfaces. *Invent. Math.*, 69(2):269–291, 1982.
- [28] Joachim Lohkamp. Discontinuity of geometric expansions. *Comment. Math. Helv.*, 71(2):213–228, 1996.
- [29] John Lott. Some geometric properties of the Bakry-Émery-Ricci tensor. *Comment. Math. Helv.*, 78(4):865–883, 2003.

- [30] John Lott. Optimal transport and Ricci curvature for metric-measure spaces. In *Surveys in differential geometry. Vol. XI*, volume 11 of *Surv. Differ. Geom.*, pages 229–257. Int. Press, Somerville, MA, 2007.
- [31] John Lott and Cédric Villani. Ricci curvature for metric-measure spaces via optimal transport. *Ann. of Math. (2)*, 169(3):903–991, 2009.
- [32] Zhiqin Lu and Julie Rowlett. Eigenvalues of collapsing domains and drift Laplacians. *Math. Res. Lett.*, 19(3):627–648, 2012.
- [33] Li Ma and Sheng-Hua Du. Extension of Reilly formula with applications to eigenvalue estimates for drifting Laplacians. *C. R. Math. Acad. Sci. Paris*, 348(21-22):1203–1206, 2010.
- [34] Emanuel Milman. On the role of convexity in isoperimetry, spectral gap and concentration. *Invent. Math.*, 177(1):1–43, 2009.
- [35] Frank Morgan. Manifolds with density. *Notices Amer. Math. Soc.*, 52(8):853–858, 2005.
- [36] Frank Morgan. Manifolds with density and Perelman’s proof of the Poincaré conjecture. *Amer. Math. Monthly*, 116(2):134–142, 2009.
- [37] Ovidiu Munteanu and Jiaping Wang. Smooth metric measure spaces with non-negative curvature. *Comm. Anal. Geom.*, 19(3):451–486, 2011.
- [38] Ovidiu Munteanu and Jiaping Wang. Analysis of weighted Laplacian and applications to Ricci solitons. *Comm. Anal. Geom.*, 20(1):55–94, 2012.
- [39] Ovidiu Munteanu and Mu-Tao Wang. The curvature of gradient Ricci solitons. *Math. Res. Lett.*, 18(6):1051–1069, 2011.
- [40] N. Nadirashvili. Berger’s isoperimetric problem and minimal immersions of surfaces. *Geom. Funct. Anal.*, 6(5):877–897, 1996.
- [41] Alberto G. Setti. Eigenvalue estimates for the weighted Laplacian on a Riemannian manifold. *Rend. Sem. Mat. Univ. Padova*, 100:27–55, 1998.
- [42] Junya Takahashi. The gap of the eigenvalues for p -forms and harmonic p -forms of constant length. *J. Geom. Phys.*, 54(4):476–484, 2005.
- [43] Guofang Wei and Will Wylie. Comparison geometry for the Bakry-Emery Ricci tensor. *J. Differential Geom.*, 83(2):377–405, 2009.
- [44] Jia-Yong Wu. Upper bounds on the first eigenvalue for a diffusion operator via Bakry-Émery Ricci curvature. *J. Math. Anal. Appl.*, 361(1):10–18, 2010.
- [45] Jia-Yong Wu. Upper Bounds on the First Eigenvalue for a Diffusion Operator via Bakry-Émery Ricci Curvature II. *Results Math.*, 63(3-4):1079–1094, 2013.

UNIVERSITÉ DE NEUCHÂTEL, LABORATOIRE DE MATHÉMATIQUES, 13 RUE E. ARGAND,
2007 NEUCHÂTEL, SWITZERLAND.

E-mail address: Bruno.Colbois@unine.ch

UNIVERSITÉ FRANÇOIS RABELAIS DE TOURS, CNRS, LABORATOIRE DE MATHÉMATIQUES
ET PHYSIQUE THÉORIQUE, UMR 7350, PARC DE GRANDMONT, 37200 TOURS, FRANCE.

E-mail address: elsoufi@univ-tours.fr

DIPARTIMENTO DI METODI E MODELLI MATEMATICI, SAPIENZA UNIVERSITÀ DI ROMA,
VIA ANTONIO SCARPA 16, 00161 ROMA, ITALY

E-mail address: savo@dmmm.uniroma1.it